

# A Rare Case of Rheumatoid Tenosynovitis with Unilateral R.A. Mimicking Tuberculosis Tenosynovitis

Kushwanth E, Arvind Natarajan K\*, Mervinrosario PM

Department of Orthopaedics, Sree Balaji Medical College, Bharath Institute of Higher Education and Research, Chromepet, Chennai, Tamilnadu, India

## ABSTRACT

Rheumatoid arthritis is an autoimmune disease chronic in nature targeting multiple joints. Savithri 48 years old female hailing from Keelkatalai house maid by profession came to Sree Balaji Medical College and Hospital in the orthopaedic OPD. This patient with atypical asymmetrical unilateral multiple joint shows a rare presentation of rheumatoid arthritis with flexor tendon synovitis shows the presence of atypical presentations and the importance of obtaining histopathological tissue diagnosis.

**Key words:** Tenosynovitis, MRI, Arthritis

**HOW TO CITE THIS ARTICLE:** Kushwanth E, Arvind Natarajan K, Mervinrosario PM, A Rare Case of Rheumatoid Tenosynovitis with Unilateral R.A. Mimicking Tuberculosis Tenosynovitis, J Res Med Dent Sci, 2022, 10 (11): 153-155.

**Corresponding author:** Dr. Arvind Natarajan K

**E-mail:** arvindmsortho@gmail.com

**Received:** 02-Sep-2022, Manuscript No. JRMDS-22-57342;

**Editor assigned:** 06-Sep-2022, PreQC No. JRMDS-22-57342 (PQ);

**Reviewed:** 21-Sep-2022, QC No. JRMDS-22-57342;

**Revised:** 03-Nov-2022, Manuscript No. JRMDS-22-57342 (R);

**Published:** 10-Nov-2022

## INTRODUCTION

Rheumatoid arthritis is an autoimmune disease chronic in nature targeting multiple joints. The synovium is involved as the first site of the inflammatory process, which if not addressed might lead to irreversible damage to the nearby cartilage and bone [1]. It is now known that autoantibodies that are characteristic of RA, and further tests such as anti CCP, RF are present positive even before the clinical presentation. Once rheumatoid arthritis is established, a number of mechanisms serve which leads to the persistence of the disease [2,3]

A proper detailed review of the mechanisms that initiate and proceed for therapeutic and prevention strategies. Rheumatoid arthritis affects the flexor tendons of the finger in about 25-40% individuals but goes undetected due to no symptoms and further involvement over the flexor tendon at the forearm tendons are very rare, tuberculosis tenosynovitis has same symptoms which mimics the problem [4].

The transition from a normal to nonspecific inflammatory synovium and subsequently to an established rheumatoid arthritis synovium is the result of a breakdown of immune tolerance both at the systemic level and at the synovial tissue level. The presence of circulating ACPA and RF, years before the appearance of clinical symptoms and the first

histological changes in the (presumed) unaffected joints of patients with RA indicates that the first stages of this degradation of immune tolerance are subclinical [5]. The importance of mediators of innate immunity in initiating synovial responses is increasingly recognized. This review will describe the main histopathology and immune histologic features of RA synovitis and address some potentially important mechanisms that may facilitate the initiation and progression of rheumatoid arthritis synovitis.

## CASE PRESENTATION

Savithri 48 years old female hailing from keelkatalai house maid by profession came to Sree Balaji Medical College and Hospital in the orthopaedic OPD. With complains of swelling over the right wrist and ankle since 3 years. Patient also complained of pain in flexing the fingers over the right hand since 1 year. Swelling over the wrist was 8\*5 CMS and was soft in consistency. Pain was insidious in onset dull aching more in flexion. Patient complains of low back ache since 6 months which was no radiating. There was no history of morning stiffness; patient gives no history of episodic increased pain with fever in the evening. No history of loss of appetite and loss of weight. Patient gives no other history of any co-morbidity in the past. With clinical examination and history patient was clinically diagnosed with tuberculosis tenosynovitis (Figures 1-4) [6].




Figure 1: MRI report of wrist and hand.



Figure 2: Clinical picture and operative picture.




Figure 3: MRI report of leg and ankle.




Figure 4: Histopathology report.

Patient was worked up for the same with chest x-ray, mantoux and other basic investigations. Where only ESR and CRP were raised and other modalities. Patient was investigated for MRI of the foot and ankle suggestive of infective/inflammatory tenosynovitis. On further rheumatologically work up was done in which anti CCP was raised taking the diagnosis in favour of Rheumatoid arthritis with tenosynovitis. Later when x-ray of bilateral hand and foot, and spine x-ray was taken which showed arthritic features of unilateral involvement of the SI joint, PIP of right hand and PIP of the right foot [7-10].

Patient was further taken for diagnostic and therapeutic total synovectomy was done in which had no rice body granuloma; rather it was greyish white with rubbery consistency.

On histopathology the features suggested for rheumatoid involvement of the synovial sheath covering the flexor tendon.

### DISCUSSION

India having a burden of tuberculosis all throughout, synovitis with even rise in temperature leads to a focused diagnosis of tuberculosis. This patient with atypical asymmetrical unilateral multiple joint shows a rare presentation of rheumatoid arthritis with flexor tendon synovitis shows the presence of atypical presentations and the importance of obtaining histopathological tissue diagnosis. Importance of serological assays like anti CCP to reach an accurate diagnosis and to provide optimum treatment to the patients.

### CONCLUSION

Patient was prescribed with DMARDS and was started with physiotherapy, after which on follow up after 3 months, the pain over the hip and fingers subsided and patient there was no decrease in the swelling of the ankle, later patient was operated for synovectomy over the right ankle after a total period of 4 months from presentation. Patient is now comfortable with taking the medication and after 9 months of initial presentation patient regained full range of motion without pain.

### REFERENCES

1. Rantapaa Dahlqvist S, De Jong BA, Berglin E, et al. Antibodies against cyclic citrullinated peptide and IgA rheumatoid factor predict the development of rheumatoid arthritis. *Arthritis Rheum* 2003; 48:2741-2749.
2. Weinstein SL, Buckwalter JA. Turek's Orthopaedics. Principles and their application. Lippincott Williams and Wilkins, 2005; 30.
3. Solomon L, Warwick D, Nayagam S, et al. Apley's system of orthopaedics and fractures. 9<sup>th</sup> edition, CRC press. *J Malays Orthop* 2010; 4.
4. Hong SH, Kim SM, Ahn JM, et al. Tuberculosis versus pyogenic arthritis: MR imaging evaluation. *Radiol* 2001; 218:848-853.
5. Kraan MC, Versendaal H, Jonker M, et al. Asymptomatic synovitis precedes clinically manifest arthritis. *Arthritis Rheum* 1998; 41:1481-1488.
6. Erickson AR, Cannella AC, Mikuls TR. clinical features of rheumatoid arthritis. 10<sup>th</sup> edition, Kelley and Firestein's *Textbook of Rheumatology*. 2017; 2:1167-1186.
7. Dyer SM, Simmons BP. The wrist and hand. *Rheumatol* 2015.

8. Sweeney SE, Harris ED, Firestein GS. Clinical features of Rheumatoid arthritis. Kelley's textbook of rheumatology. 2013.
9. Simmen BR, Kolling C, Herren DB. Management of rheumatoid wrist. Curr Orthopract 2007; 21:344-357.
10. Saffar P. Flexor tendon synovectomy in rheumatoid arthritis. Int Congr Ser 2006; 1295:107-117.