

# Changes in Alveolar Bone Dimension after Extraction Sockets and Methods of ridge preservation

Saumya Jitendra Kanode\*, Anand Narayanrao Wankhede

Department of Periodontology, Sharad Pawar Dental College and Hospital, Datta Meghe Institute of Medical Sciences, Sawangi, Wardha, Maharashtra, India

## ABSTRACT

The alveolar process develops during tooth formation and goes through bone atrophy after a tooth or teeth are lost. This could happen because of periodontal disease, periapical pathologies, and mechanical trauma which cause loss of surrounding bone around the tooth. Thus, leads to the resorption of the alveolar ridge. Also, predisposing factors that affect alveolar bone loss following tooth extraction such as age, gender, systemic conditions, facial morphology, and functional stress on the extraction wound.

Post extraction complications include loss of function, a reduction in vertical height, horizontal width, insufficient bone for dental implants, and prosthodontics difficulties. It has been noted that the size of the residual ridge shrinks more quickly within the first six months. After extraction, the maxillary arch tended to have more horizontal resorption together with vertical bone resorption than the mandible. In order to achieve successful implant results, both the bone and soft tissue contour should be preserved as well as possible.

Alveolar Ridge Preservation (ARP) is considered as a Guided Bone Regeneration (GBR) technique for preventing ridge resorption succeeding tooth extraction. The aim is to provide the existing research on alveolar bone loss and tissue alterations in extraction sockets, the effects of age on bone resorption, and different Ridge preservation strategies.

**Key words:** Alveolar Bone, Extraction sockets, Ridge preservation, Methods

**HOW TO CITE THIS ARTICLE:** Saumya Jitendra Kanode, Anand Narayanrao Wankhede, Changes in Alveolar Bone Dimension after Extraction Sockets and Methods of ridge preservation, J Res Med Dent Sci, 2022, 10 (11): 000-000

**Corresponding author:** Saumya Jitendra Kanode

**E-mail:** kanodesaumy@gmail.com

**Received:** 01-Sep-2022, Manuscript No. JRMDS-22-49475;

**Editor assigned:** 05-Sep-2022, PreQC No. JRMDS-22-49475 (PQ);

**Reviewed:** 19-Sep-2022, QC No. JRMDS-22-49475;

**Revised:** 02-Nov-2022, Manuscript No. JRMDS-22-49475 (R);

**Published:** 09-Nov-2022

## INTRODUCTION

Resorption of the alveolar bone varies among individuals to individuals, sites to sites after extraction which leads to loss of function, insufficient bone for dental implants, and prosthodontics difficulties [1]. Resorption of mandibular ridge height is more than maxillary ridge height, while buccal ridge width loss is greater [2,3].

One of the most common dental treatments that result in the reduction of vertical ridge height is tooth extraction. This is particularly noticeable on the buccal aspect, according to reports [4-6]. This resorption process results in a narrower and shorter ridge [7]. The ridge is shifted to a more palatal/lingual position as a consequence of constriction of the blood clot and the existence of bundle bone at the crystal region. It has been also seen that periodontitis, periapical lesion, or trauma to the bone and teeth may have caused alveolar bone loss prior to tooth extraction [6-13].

The size of the residual ridge shrinks most quickly in the first six months, although resorption of bone in the residual ridge continues at a slower pace throughout life, resulting in depletion in the horizontal and vertical width of the jaw structure [13,14].

Extraction sites had a greater proportion of vital bone in the maxilla when they healed, whereas preservation sites had identical percentages of vital bone in the mandible and maxilla when they healed.

In general, after tooth loss, alveolar bone remodelling results in smaller alveolar ridge extent in horizontal and vertical planes. According to studies, ridge changes caused by tooth loss can cause a 40% to 60% bone loss in vertical and horizontal dimensions in as little as 3 months [14,15].

## LITERATURE REVIEW

**Age factor for alveolar bone:** Amler, et al. looked at the role of age in wound healing after extraction. Individuals in their second decennium of life were in contrast to those in their sixth decennium or above in terms of healing time. After the first ten days of the post-extraction phase, no significant variance in the rate of healing between young and elderly individuals were observed; however, after ten days, the rate of tissue regeneration in younger individuals

began to accelerate as compared to older patients. In the same finding is seen after 20 days the rate of tissue regeneration began to accelerate. After 30 days, both young and elderly people had healed at the same rate. The length of active revitalization in younger and older people is dependent on the timing of donor [15,16].

Subjects over 50 years old had more horizontal and vertical resorption after extraction than those under the age of 50. In the age less than of 50 years, the ridge preservation approach compensated for the decrease in vertical ridge height, but not in the ≥ 50 years of age. The effect was similar for both age groups, although the ridge preservation effect was similar for both the age group [14,15].

**Changes seen in tissue morphology of extraction sockets in humans**

The ridge changes in tandem accompanied by the recovery of the soft as well as hard tissue wounds, but the remodelling procedure can carry on even after *de novo* bone creation in the socket has stopped [16,17].

Cardaropoli, et al. in his study he looked at how fresh extraction sockets healed in the coronal, central, and apical compartments over a 6 month period. The scientists found that the socket was pervaded with the woven bone after 4 weeks after recovery and that this immature bone was restored with lamellar bone and marrow after 2 months. The buccal wall of the socket had significant resorption at the same time as these intra alveolar healing activities. As a result, the height of the buccal wall of the socket was reduced by 2.5 mm on average when compared to the lingual wall [17,18].

Tissue generation following tooth extraction was also studied in human models. He came to the conclusion that the blood clot that had previously pervaded the socket's entry had been replaced with granulation tissue. Osteoid

production began after one week of tissue modelling, and after about six weeks, the socket's periphery included islands of immature woven bone [18,19].

Johnson was the first to show that tooth extraction could result in a decrement in height and width of 2.5 to 7 mm and up to 3 mm. moreover, they came across that the majority of the variations happened in the first month, with a little extra drop in the ridge persisting for periods varies from 10 to 20 weeks [19,20].

Petrokovski and Massler reached a similar conclusion, they noted that tissue devitalisation occurs more in the buccal wall of the molar area than in the aesthetic zone [20]. Schropp, et al. stated that a year after tooth extraction, the remnant alveolar ridge can shrink by up to 50% in width [16,17].

They theorized that because the bundle bone is part of the periodontium, removing a tooth leaves it useless, and resorption occur [6]. According to other researchers, the collapse of the buccal soft tissue resulted in significant buccal oral changes. Other authors, on the other hand, have studied that surgical trauma during tooth extraction may result in the periosteum's separation from the underlying bone surface. This could result in injury to the vessels and an initial inflammatory reaction, which will then mediate bone tissue resorption [21-25].

**DISCUSSION**

**Various ridge preservation modalities**

Numerous ridge preservation approaches have been proposed to influence bone remodelling after extraction. Routinely, Ridge preservation is described as a method performed at the time of tooth extraction with the goal of reducing exterior ridge resorption and optimizing bone growth within the socket [26-35].

**Table 1:** Shows various ridge preservation modalities.

Sr.no	Author (Year)	Title	Conclusion of study
1	Lauren A Brownfield, et al.	Alveolar Ridge Preservation (ARP) is a Guided Bone Regeneration (GBR) procedure aiming to control post-extraction ridge resorption.	No statistically significant differences were found in dimensional changes between groups after 10 to 12 weeks of healings. Bone loss in the vertical height was less likely to occur when the average buccal plate thickness was 1.3 mm
2	Daniele Cardaropoli, et al.	Socket preservation using bovine mineral and collagen	Socket preservation using bone material(bovine) and collagen membrane(porcine) considerably limits the amount of horizontal as well as vertical bone resorption when compared with the extraction of tooth alone
3	HammerleCH Schmid, et al.	A novel model system for the study of experimental guided bone regeneration in humans	The presented model system is suitable to study tissue physiology of bone regeneration in humans with minimal complications or adverse effects to the volunteers.
4	Lanka Mahesh, et al.	Alveolar ridge preservation with the socket plug technique utilizing an alloplastic bone substitute (Putty) or a particulate xenograft: A histological pilot study.	The amount of new bone regenerated was also statistically significantly greater in the alloplasts study group as compared to the xenograft group. Results suggest that ridge preservation using alloplastic (calcium phosphosilicate) bone substitute showed more timely graft substitution and increased bone

			regeneration when compared to a bovine bone xenograft.
5	Eric Todd Scheyer, et al.	A randomized, controlled, multicentre clinical trial of post-extraction alveolar ridge preservation	Treatments were demineralized allograft plus reconstituted and cross-linked collagen membrane (DFDBA)+(RECCX) or deproteinized bovine bone mineral with collagen plus native, bilayer collagen membrane (DBBMC+NBCM). DBBMC +NBCM provided better soft tissue healing and ridge preservation for implant placement.
6	G Avila-Ortiz, et al.	Efficacy of alveolar ridge preservation RCT.	The Control group includes tooth extraction and the experimental group received Alveolar Ridge Preservation using a combination particulate bone allograft with a non-absorbable membrane (dPTFE) following tooth extraction. Bone resorption was significantly greater in the control group. No significant differences with respect to soft tissue contour change were observed between groups.
7	Veronica J Lai, et al.	Ridge preservation following tooth extraction using xenograft (bovine) compared with xenograft (porcine)-RCT	Ridge preservation with the help of xenograft (porcine) results in dimensional stability with xenograft(bovine).
8	Yiping Wei, et al.	Ridge preservation in maxillary molar extraction sites with severe periodontitis: A prospective observational clinical trial	Ridge preservation can improve alveolar ridge dimensions and decrease the necessity of advanced regenerative procedures at implant placement compared to natural healing in the maxillary molar extraction sockets with severe periodontitis.
9	Brend Pjotr Jonker, et al.	Soft tissue contour and radiographic evaluation of ridge preservation in early implant placement: A randomized controlled clinical trial	Ridge preservation using a bone substitute (xenogenic) covered with a collagen matrix or a palatal graft, results in less bone resorption.

**CONCLUSION**

Studies have shown that by using various graft materials and techniques, Methods of ridge preservation limit the resorption of alveolar ridge width and provide gain in hard tissue ridge height when compared to extraction alone. Thus, it limits bone resorption and helps in the successful results of dental implant placement

**REFERENCE**

1. Bartee BK. Extraction site reconstruction for alveolar ridge preservation. Part 2: membrane-assisted surgical technique. J Oral Implantol 2001; 27:194-197.
2. Smukler H, Landi L, Setayesh R. Histomorphometric evaluation of extraction sockets and deficient alveolar ridges treated with allograft and barrier membrane: A pilot study. Int J Oral Maxillofac Implants 1999; 14:407-416.
3. Iasella JM, Greenwell H, Miller RL, et al. Ridge preservation with freeze dried bone allograft and a collagen membrane compared to extraction alone for implant site development: A clinical and histologic study in humans. J Periodontol 2003; 74:990-999.
4. Lekovic V, Kenney EB, Weinlaender M, et al. A bone regenerative approach to alveolar ridge

maintenance following tooth extraction. Report of 10 cases. J Periodontol 1997; 68:563-570.

5. Lekovic V, Camargo PM, Klokkevold PR, et al. Preservation of alveolar bone in extraction sockets using bioabsorbable membranes. J Periodontol 1998; 69:1044-1049.
6. Araujo MG, Lindhe J. Dimensional ridge alterations following tooth extraction. An experimental study in the dog. J Clin Periodontol 2005; 32:212-218.
7. Pinho MN, Roriz VL, Novaes AB Jr, et al. Titanium membranes in prevention of alveolar collapse after tooth extraction. Implant Dent 2006; 15:53-61.
8. Wankhede AN, Dhadse PV, Jaiswal PG, et al. Comparative evaluation of interleukin-17 in gingival crevicular fluid of patients with aggressive periodontitis and healthy gingival sites. J Datta Meghe Inst Med Sci Univ 2021; 16:90-93.
9. Misch CE, Dietsch-Misch F, Misch CM. A modified socket seal surgery with composite graft approach. J Oral Implantol 1999; 25:244-250.
10. Jain P, Kumar CMR, Jain M, et al. A split mouth study on crystal bone architectural changes by flapped and flapless implant surgical procedures. J Pharma Res Intl 2021; 33:103-111.
11. Nambiar KM, Kamble R, Suroliya KP, et al. Alveolar Bone Grafting-A Summary of History and Current

- Literature on the Technique. J Evol Med Dent Sci 2020; 9:2652-2657.
12. Ghoshal PK, Kamble RH, Shrivastav SS, et al. Radiographic evaluation of alveolar bone dimensions in the inter-radicular area between maxillary central incisors as "safe zone" for the placement of miniscrew implants in different growth patterns a digital volume tomographical study. J Evo Med Dent Sci 2019; 8:3836-3840.
  13. Pietrokovski J, Starinsky R, Arensburg B, et al. Morphologic characteristics of bony edentulous jaws. J Prosthodont 2007; 16:141-147.
  14. Jahangiri L, Devlin H, Ting K, et al. Current perspectives in residual ridge remodelling and its clinical implications: a review. J Prosthet Dent 1998; 80:224-237.
  15. Barone A, Ricci M, Tonelli P, et al. Tissue changes of extraction sockets in humans: A comparison of spontaneous healing vs. ridge preservation with secondary soft tissue healing. Clin Oral Impl Res 2012; 1-7.
  16. Amler MH. Age factor in human alveolar bone repair. J Oral Implantol 1993; 19:138-142.
  17. Schropp L, Wenzel A, Kostopoulos L, et al. Bone healing and soft tissue contour changes following single-tooth extraction: a clinical and radiographic 12 month prospective study. Int J Periodontics Restorative Dent 2003; 23:313-323.
  18. Cardaropoli G, Araujo M, Hayacibara R, et al. Healing of extraction sockets and surgically produced-augmented and non-augmented-defects in the alveolar ridge. An experimental study in the dog. J Clin Periodontol 2005; 32:435-440.
  19. Evian CI, Rosenberg ES, Coslet JG, et al. The osteogenic activity of bone removed from healing extraction sockets in humans. J Periodontol 1982; 53:81-85.
  20. Johnson K. A study of the dimensional changes occurring in the maxilla following tooth extraction. Aust Dent J 1969; 14:241-244.
  21. Pietrokovski J, Massler M. Alveolar ridge resorption following tooth extraction. J Prosthet Dent 1967; 17:21-27.
  22. Wilderman, Malbern N. "Repair after a Periosteal Retention Procedure." J Periodontol 34:487-503.
  23. Staffileno H, Levy S, Gargiulo A. Histologic study of cellular mobilization and repair following a periosteal retention operation *via* split thickness mucogingival flap surgery. J Periodontol 1966; 37:117-131.
  24. Wood DL, Hoag PM, Donnenfeld OW, et al. Alveolar crest reduction following full and partial thickness flaps. J Periodontol 1972; 43:141-144.
  25. Bragger U, Pasquali L, Kornman KS. Remodelling of interdental alveolar bone after periodontal flap procedures assessed by means of Computer Assisted Densitometry Image Analysis (CADIA). J Clin Periodontol 1988; 15:558-564.
  26. Durge KJ, Baliga VS, Sridhar SB, et al. Extraction socket grafting using recombinant human bone morphogenetic protein 2 clinical implications and histological observations. BMC Res Notes 2021; 14:61.
  27. Brownfield LA, Weltman RL. Ridge preservation with or without an osteoinductive allograft: A clinical, radiographic, micro computed tomography, and histologic study evaluating dimensional changes and new bone formation of the alveolar ridge. J Periodontol 2012; 83:581-589.
  28. Cardaropoli D, Tamagnone L, Roffredo A, et al. Socket preservation using bovine bone mineral and collagen membrane: a randomized controlled clinical trial with histologic analysis. Int J Periodontics Restorative Dent 2012; 32:421-430.
  29. Hammerle CH, Schmid J, Olah AJ, et al. A novel model system for the study of experimental guided bone formation in humans. Clin Oral Implants Res 1996; 7:38-47.
  30. Mahesh L, Venkataraman N, Shukla S, et al. Alveolar ridge preservation with the socket-plug technique utilizing an alloplastic putty bone substitute or a particulate xenograft: a histological pilot study. J Oral Implantol 2015; 41:178-183.
  31. Scheyer ET, Heard R, Janakievski J, et al. A randomized, controlled, multicentre clinical trial of post extraction alveolar ridge preservation. J Clin Periodontol 2016; 43:1188-1199.
  32. Avila-Ortiz G, Gubler M, Romero-Bustillos M, et al. Efficacy of alveolar ridge preservation: a randomized controlled trial. J dent res 2020; 99:402-409.
  33. Lai VJ, Michalek JE, Liu Q, et al. Ridge preservation following tooth extraction using bovine xenograft compared with porcine xenograft: A randomized controlled clinical trial. J periodontol 2020; 91:361-368.
  34. Wei Y, Xu T, Zhao L, et al. Ridge preservation in maxillary molar extraction sites with severe periodontitis: A prospective observational clinical trial. Clin Oral Investig 2021.
  35. Jonker BP, Gil A, Naenni N, et al. Soft tissue contour and radiographic evaluation of ridge preservation in early implant placement: A randomized controlled clinical trial. Clin Oral Implants Res 2021; 32:123-133.