

# Evaluation of Photodynamic Therapy in Comparison with Conventional Treatment for the Treatment of Periodontal Disease-A Clinical Trial

Iram Rafique Pawane, Jaiganesh Ramamurthy\*

Department of Periodontics, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences (SIMATS) Saveetha University, Chennai, India

# ABSTRACT

Aim: The aim of this study was to compare the clinical results of conventional non-surgical periodontal therapy with or without adjunct PDT in patients of chronic periodontitis.

Materials and methods: 20 patients with chronic untreated periodontitis, (8 female, 12 male, mean aged 36.35 years, all non-smokers, Systemic diseases, including anti-inflammatory, blood stimulants, or systemic antibiotics, within the last 6 months were exempted from the criteria, previously untreated chronic periodontitis; in any quadrant at least 1 premolar and 1 molar with at least 4 teeth each; in each quadrant at least 1 tooth with an attachment loss of>3mm were included in the study. The periodontal status of each subject was evaluated at baseline and 6 weeks following periodontal therapy. PDs, clinical attachment level (CAL) were assessed.

All patients received non-surgical periodontal treatment comprising a thorough scaling and root planning of all periodontal involved teeth using a split mouth design, two quadrants were additionally treated with PDT, with a designated photosensitizer dye (methylene blue) and a diode laser (biolase 940nm). Laser application at six locations per tooth was carried out circumferentially.

The subjects were given thorough oral hygiene maintenance instructions and recalled after two weeks for a second round of PDT at the same sites. A total of two exposures for the photodynamic therapy group with a two week interval were done Probing depths and attachment levels were assessed after 6 weeks. Statistical analysis was done to compare intergroup using the paired t test. Differences were considered as statistically significant at p<0.05.

Results: Baseline mean values for PD and CAL were significantly different in the test and control group. Values for CAL and PD decreased significantly 6 weeks after treatment in the control group, with a higher impact on the sites treated with adjunctive PDT

*Conclusion: In patients with untreated chronic periodontitis, clinical outcomes of conventional non-surgical periodontal therapy can be improved by multiple PDT.* 

Key words: Diode laser, Photodynamic therapy, Non-surgical periodontal therapy

**HOW TO CITE THIS ARTICLE:** Iram Rafique Pawane, Jaiganesh Ramamurthy, Evaluation of Photodynamic Therapy in Comparison with Conventional Treatment for the Treatment of Periodontal Disease-A Clinical Trial, J Res Med Dent Sci, 2022, 10(1): 513-518

Corresponding author: Iram Rafique Pawane E-mail⊠:iramrafique2396@gmail.com Received: 24/10/2021 Accepted: 31/12/2021

# INTRODUCTION

Periodontitis is caused by inflammation of the supporting teeth structure and by various periodontopathic bacteria in response to chronic infections, respectively. Current concepts are based on mechanical scaling and root planning to remove bacterial deposits, cementum contaminated with the bacteria and endotoxins in the treatment of periodontal involved teeth. However in sites where the mechanical scaling and root planning are difficult to access the removal of plaque and the reduction of the number of infectious cells. In cases which do not respond to conventional treatments, certain therapeutic alternatives, such as systemic and local antibiotics have recently gained popularity as a possible replacement or

complement to traditional mechanical debridement. CO2 been used [1,2]. Laser-assisted periodontal therapy has laser, neodymium doped: yttrium-aluminum-garnet (Nd:YAG) laser, and diode and erbium-doped: yttriumaluminum-garnet (ER:YAG) laser (Er:YAG) Laser have been used in periodontal pocket therapy for both hard and soft tissue treatment.

During irradiation, some of the laser energy scatters and penetrates into periodontal pockets. The low-energy attenuated laser can then stimulate the cells of the surrounding tissue, reducing inflammatory conditions in cell proliferation and increasing lymph flow, improving periodontal tissue attachment and probably reducing postoperative pain [3,4].

The wavelength of a diode laser is 810 nm or 910–980 nm, which has no effect on dental hard tissues. As a result, the laser is an ideal soft tissue surgical laser that can be used to cut and coagulate gingiva and oral mucosa, as well as for

soft tissue curettage and sulcular debridement. It has a bactericidal effect as well [4,5].

PDT is the light energy process used to activate a photo sensitizer in the presence of oxygen. The principle of operation is that the photosensitizer undergoes a transition to a higher energy condition that produces highly reactive oxygen conditions [6]. This oxygen alone could cause microorganisms toxicity. A number of photosenses against target microorganisms have proved to be effective without damaging the host tissues. PDT can be effective in killing bacteria like Porphyromonas gingivalis or Fusobacterium nucleatum in vitro which periodontopathogens [7] are. The photo sensitization of *P. gingivalis* was demonstrated by using an animal model in vivo, which resulted in a decrease in bone loss [8]. In a model of the Beagle dog, inflammatory signs were positive and *P. gingivalis* could be suppressed [9]. Similar clinical results have been shown in the evaluation of PDT's impact on the treatment of human aggressive periodontitis, photosensitisation and SRP [10-12]. The comparison of conventional debridement with or without the additional use of PDT in chronic periodontitis showed improved clinical parameter improvement in the PDT group [13,14].

This study was intended to compare the clinical results of conventional non-surgical periodontal therapy with or without adjunct PDT in patients of chronic periodontitis by testing for non-surgical periodontal therapy hypothesizing that incorporating that PDT as an adjunct could improve clinical outcomes

# MATERIALS AND METHODS

## Patient selection

20 patients, each of whom presented chronic untreated periodontitis, were recruited from the outpatient Department of Periodontics and Oral Implantology, Saveetha Institute of Medical and Technical Sciences, Chennai (8 female, 12 male, mean aged: 36.35 years, all non-smokers).

## **Exclusion criteria**

Systemic diseases, including anti-inflammatory, blood stimulants, or systemic antibiotics, could influence periodontal therapy outcomes within the last 6 months were exempted from the criteria.

## Inclusion criteria

The study included the following: previously untreated chronic periodontitis; in any quadrant at least 1 premolar and 1 molar with at least 4 teeth each; in each quadrant at least 1 tooth with an attachment loss of >3mm.

#### **Informed consent**

All patients were informed of the study and a written and video consent was taken prior to the trial in the period from December 2020 to February 2021, to participate in the study for 6 Weeks. The study was carried out in full

conformity with the declared ethical principles approved by the Ethics Committee of the University.

## **Clinical parameters**

The periodontal status of each subject was evaluated at baseline and 6 weeks following periodontal therapy. A blind examiner, who was not involved in the study of patients, documented the PDs, relative attachment level (RAL), degree of tooth mobility, and furcation involvement. An experienced periodontal examiner conducted all measurements allowing intra-experimental value comparison.

The upper and lower teeth have been recorded by taking impressions followed by the fabrication of personalised splints fitting with the teeth. These splits were used to provide both PD and relative attachment status with reproducible measurement points. Thus, the individual splints were produced by a vacuum-forming process for each subject. Only short of tooth emergence was trimmed. A groove was created in the splint for each studied site and a line for the calibrating pressure periodontal probe was formed to facilitate a reproducible sensor position during measurements.

# **Treatment procedure**

All patients received non-surgical periodontal treatment comprising a thorough scaling and root planning of all periodontal involved teeth employing both hand.

Instruments (Gracey curettes, Hu-Friedy, Leimen, Germany) and a piezo-electric ultrasonic hand-piece with a slim-line styled scaler tip (Woodpecker) by the same clinician. Using a split-mouth design, two quadrants (test group) were additionally treated with PDT. Therefore, after periodontal debridement, the quadrants were assigned to different groups according to a randomised lot picking technique. The sequence was concealed until interventions were assigned.

In combination with a designated photosensitizer dye (methylene blue) the PDT was performed by a diode laser (biolase 940nm). Periodontal pockets have been rinsed from the bottom of the pocket using the photosensitizer and the blunt cannula to complete the pocket filling and roots coating. The pockets were rinsed with water for excess photosensitizer after 3minutes of residence time. The remaining photo sensor was activated for 10s per site using the laser probe.

Laser application at six locations per tooth was carried out circumferentially. The subjects were given thorough oral hygiene maintenance instructions and recalled after two weeks for a second round of PDT at the same sites. A total of two exposures for the photodynamic therapy group with a two week interval were done Probing depths and attachment levels were assessed after 6 weeks.

# Statistical analysis

For statistical analysis IBM SPSS version 23 (SPSS Inc., Chicago, IL, USA). Comparison within the groups with

respect to the treatment intervals were performed using the paired t test. Differences were considered as statistically significant at p < 0.05.

## RESULTS

# **Probing depths**

Baseline PDs of periodontal involved teeth were statistically significant in the test group and control group (p<0.05). Six weeks after treatment, in both groups, a decrease in PDs could be found (p<0.05) with a higher impact in the test group than in the control, with a

higher impact on the sites treated with adjunctive PDT (p<0.05).

# **Clinical attachment level**

The attachment levels of periodontal involved teeth did not differ significantly in the test group and control group at baseline (p<0.05). After 6 weeks, a lower attachment gain could be observed in the control group than in the test group (p<0.05). Comparing the differences in CAL, an attachment gain could be observed in both groups, with a higher impact on the sites treated with adjunctive PDT (p<0.05).

# Table 1: Depicts statistical analysis for pre and post intervention for adjunctive photodynamic therapy in both groups; with and without photodynamic therapy.

		Mean	Significance
Group 1	PD before	$6.70 \pm 1.081$	0
	PD after	$4.55 \pm 1.146$	
	CAL before	8.30 ± 1.218	0
	CAL after	6.65 ± 1.137	0
Group 2	PD before	$6.92 \pm 1.112$	0
	PD after	$5.95 \pm 1.192$	0
	CAL before	9.12 ± 1.256	0
	CAL after	8.46 ± 1.127	0

Group 1: With adjunctive photodynamic therapy, Group 2: Without adjunctive pdt, pd: probing depth, cal: clinical attachment level.



Figure 1: Bar graph represents the association between probing depths before and after intervention in both groups; with and without adjunctive photodynamic therapy (p<0.05) group 1 depicts significant improvement in probing depth reduction than group 2 (Group 1: With photodynamic therapy group 2: without photodynamic therapy).



Figure 2: Bar graph represents the association between clinical attachment levels before and after intervention in both groups; with and without adjunctive photodynamic therapy, (p<0.05) group 1 significant improvement depicts in clinical attachment gain than group 2 (Group 1: with photodynamic Without therapy group 2: photodynamic therapy).

## DISCUSSION

This study demonstrates that PDT procedures improved the clinical results for the non-surgical periodontal treatment of chronic periodontitis. The conventional mechanical device of the root is seen as a prerequisite for success in the long term [15]. Studies could, however, indicate that an additional benefit in the treatment of chronic periodontitis may come from adjunctive treatment processes like minocycline [16,17] or laser radiation [18,19]. The Er: YAG laser is a new technique for sub-gingival debridement. The residual calculus after laser irradiation can be demonstrated to depend on the fluorescence threshold level, without removing a relevant quantity of the root cementum As to the microbiological results, Er:YAG laser, curettes, sonic and ultrasonic scalers have the same effects in chronic periodontitis patients [20]. Full mouth treatment concepts are another attempt at improved periodontal therapy to avoid early re-infection in untreated areas. Disputable data on microbiological effects of full mouth non-surgical periodontal therapy compared to the normal quadrant approach have been reported. A recent study has not confirmed any re-colonization differences after SRP for 24 hours compared to treatments during several sessions [21].

In current literature, only minor differences in treatment effects among these treatment strategies were observed among adults with chronic periodontitis [22]. A systematic antibiotic constituent in the long-term management of periodontal diseases provides significant clinical benefits from the sub-antimicrobial dose doxycycline therapy as an addition to SRP [23]. However, the use of this therapy in addition to non-surgical periodontal debridement among smokers could not be demonstrated [24]. In periodontal therapy systemic antimicrobials should be an adjunct to mechanical debridement. These results validates the search for new methods of treatment to improve conventional debridement. The limited access to plaque by topical agents and the development of antibiotic resistance make alternative strategies necessary in order to control the use of biofilms and treat periodontal diseases [25]. PDT is mediated by singlet oxygen, which affects extracellular molecules directly. Polysaccharides present also have photo damage potential within the extracellular matrix of the polymers in a bacterial biofilm. Such dual activity is not antibiotic-like and can have a major benefit from aPDT. Furthermore, it seems unlikely that singlet oxygen or free radicals will develop resistance to cytotoxic action. The PDT is equally effective against antibiotic, antibiotic, resistant and susceptible bacteria, and repeated photosensitization does not induce resistant strain selection [26]. The PDT procedure involves the laser activation of photosensitizing dye. As laser light irradiated only the test quadrants, there were no effects on bacteria in the control quadrants.

The results of this study correspond to those of the study that evaluate the effect only and in combination with conventional SRP of photo disinfection. In the group treated with SRP alone after 12 weeks, the authors have assessed 33 patients with chronic periodontitis for a clinical attachment gain of 0.36 0.35 mm. For SRP with adjunctive PDT, a gain of 0.86 0.61 mm was observed. These values are within the same range as in this study; the control group has less RAL values than the adjunctive PDT values in the group. A greater reduction in the BOP in the test group was observed in both studies. Although the values for SRP with adjunctive PDT were substantially different, the differences were minor. In a 6month follow-up, however, the effects of local drug delivery on the SRP were evaluated, PD differences were observed from 0.1 to almost 0.5 mm and smaller effects for attachment gains, although statistically different differences could be observed [27]. A study assessing the effect of a chlorhexidine sub-gingival chip could detect a 0.5 mm difference in clinical attachment gain in favour of Chlorhexidine after six months [28].

However, it remains a question whether these improvements are clinically significant. The additional PDT application to SRP of one exposure did not lead to further improvements in pocket depth reduction and attachment gain, when evaluating patients who received supportive periodontal therapy, but it led to significantly higher BOP scores than SRP alone. In patients with aggressive periodontitis, SRP was also compared to PDT alone. In a split mouth design, ten patients were treated. In both groups after 3 months a significant reduction in BOPs was observed. PD values and clinical attachment levels have also reduced after 3 months [29].

. A positive effect on attachment gain reduced PD and reduced use of metronidazole plus amoxicillin as sole therapy for periodontal treatments with mechanical debridement have recently been shown. However, the clinician should expect remaining mineralized deposits on the root surface, irrespective of the use of antibiotics or aPDT as the sole treatment system. This residual sub gingival calculus may serve as a base for bacteria and help to develop pockets and to progress periodontal disease [30]. Any viable bacteria on rough surfaces of residual calculus could serve as a source of periodontal lesion re-infection and cause periodontitis to progress.

In this study, all of the patients obtained SRP periodontal therapy with both hand instruments (curettes) and a piezo-electric hand piece for all the teeth. No difference was found in the treatment of chronic periodontitis concerning clinical outcome between ultrasonic and manual debridement. In addition, each patient was treated with both debridement procedures so that quadrants treated with split mouth design could be compared intra-experimentally. A gentle gingival sulcus probing was performed with a pressure sensitive probe, since the sampling force of 20 g was demonstrated to avoid trauma of periodontal tissues during sampling. The present study shows that the use of PDT in conjunction with non-surgical treatment results has a positive impact. Increasing the number of exposures to photodynamic therapy is a newer approach to treat non-surgically as it is inferred that this could affect the microbial count. Thus, it could be possible to improve non-surgical periodontal therapy by adding antimicrobial photodynamic treatment exposures to conventional nontherapy approaches. Furthermore, surgical the microbiological effects behind the clinical benefits observed should be studied and the decrease in microbial load to repeated exposures must be observed.

# CONCLUSION

From the above study, it can be concluded that in patients with chronic periodontitis, multiple exposures of adjunctive photodynamic therapy after conventional nonsurgical therapy, i.e. scaling and root planning improves clinical outcomes.

# REFERENCES

- 1. Polson AM, Frederick GT, Ladenheim S, et al. The production of a root surface smear layer by instrumentation and its removal by citric acid. J Periodontol 1984; 55:443-6.
- 2. Kapoor A, Malhotra R, Grover V, et al. Systemic antibiotic therapy in periodontics. Dent Res J 2012; 9:505.
- 3. Kornman KS, Page RC, Tonetti MS. The host response to the microbial challenge in periodontitis: assembling the players. Periodontol 1997; 14:33-53.
- 4. Hamblin MR, Hasan T. Photodynamic therapy: A new antimicrobial approach to infectious disease?. Photochem Photobiol Sci 2004; 3:436-50.
- 5. Wainwright M. Photodynamic antimicrobial chemotherapy (PACT). J Antimicrob Chemo 1998; 42:13-28.
- 6. Daniell MD, Hill JS. A history of photodynamic therapy. Aus New Zealand J Sur 1991; 61:340-8.
- 7. Pfitzner A, Sigusch BW, Albrecht V, et al. Killing of periodontopathogenic bacteria by photodynamic therapy. J Periodontol 2004; 75:1343-9.
- 8. Komerik N, MacRobert AJ. Photodynamic therapy as an alternative antimicrobial modality for oral infections. J Enviro Patho Toxicol Oncol 2006; 25.
- 9. Sigusch BW, Pfitzner A, Albrecht V, et al. Efficacy of photodynamic therapy on inflammatory signs and two selected periodontopathogenic species in a beagle dog model. J Periodontol 2005; 76:1100-5.
- 10. Monteiro JS, Rangel EE, de Oliveira SC, et al. Enhancement of photodynamic inactivation of planktonic cultures of Staphylococcus aureus by DMMB-AuNPs. Photodiag Photodyn Thera 2020; 31:101930.
- 11. Giroldo LM, Felipe MP, de Oliveira MA, et al. Photodynamic antimicrobial chemotherapy (PACT) with methylene blue increases membrane permeability in Candida albicans. Lasers Med Sci 2009; 24:109-12.
- 12. de Oliveira RR, Schwartz-Filho HO, Novaes Jr AB, et al. Antimicrobial photodynamic therapy in the non-surgical treatment of aggressive periodontitis: A preliminary randomized controlled clinical study. J Periodontol 2007; 78:965-73.
- 13. Krause F, Braun A, Brede O, Eberhard J, et al. Evaluation of selective calculus removal by a

fluorescence feedback-controlled Er: YAG laser in vitro. J Clin Periodontol 2007; 34:66-71.

- 14. Hohwy T, Andersen KE, Sølvsten H, et al. Allergic contact dermatitis to methyl aminolevulinate after photodynamic therapy in 9 patients. Contact Derma 2007; 57:321-3.
- 15. Greenstein G. Supragingival and subgingival irrigation: Practical application in the treatment of periodontal diseases. Compendium 1992; 13:1098.
- 16. Cortelli JR, Aquino DR, Cortelli SC, et al. A doubleblind randomized clinical trial of subgingival minocycline for chronic periodontitis. J Oral Sci 2008; 50:259-65.
- 17. Cortelli JR, Querido SM, Aquino DR, et al. Longitudinal clinical evaluation of adjunct minocycline in the treatment of chronic periodontitis. J Periodontol 2006; 77:161-6.
- 18. Qadri T, Miranda L, Tuner J, et al. The short-term effects of low-level lasers as adjunct therapy in the treatment of periodontal inflammation. J Clin Periodontol 2005; 32:714-9.
- 19. Cobb CM. Lasers in periodontics: a review of the literature. J Periodontol 2006; 77:545-64.
- 20. Derdilopoulou FV, Nonhoff J, Neumann K, et al. Microbiological findings after periodontal therapy using curettes, Er: YAG laser, sonic, and ultrasonic scalers. J Clin Periodontol 2007; 34:588-98.
- 21. Jervøe-Storm PM, AlAhdab H, Koltzscher M, et al. Comparison of curet and paper point sampling of subgingival bacteria as analyzed
- 22. by real-time polymerase chain reaction. J Peridontol 2007; 78:909-17.
- 23. Eberhard J, Jepsen S, Jervøe-Storm PM, et al. Fullmouth treatment modalities (within 24 hours) for chronic periodontitis in adults. Cochrane Database Syst Rev 2015.
- Gurkan A, Çınarcık S, Hüseyinov A. Adjunctive subantimicrobial dose doxycycline: Effect on clinical parameters and gingival crevicular fluid transforming growth factor-β1 levels in severe, generalized chronic periodontitis. J Clin Periodontol 2005; 32:244-53.
- 25. Needleman I, Suvan J, Gilthorpe MS, et al. A randomized-controlled trial of low-dose doxycycline for periodontitis in smokers. J Clin Periodontol 2007; 34:325-33.
- 26. Goslinski T, Osmalek T, Konopka K, et al. Photophysical properties and photocytotoxicity of novel phthalocyanines–potentially useful for their application in photodynamic therapy. Polyhed 2011 26; 30:1538-46.
- 27. Bonito AJ, Lux L, Lohr KN. Impact of local adjuncts to scaling and root planning in periodontal disease therapy: A systematic review. J Periodontol 2005; 76:1227-36.
- 28. Paolantonio Michele, Giuseppe Perinetti, Simonetta D'Ercole. Internal decontamination of

dental implants: an in vivo randomized microbiologic 6-month trial on the effects of a chlorhexidine gel. J Periodontol 2008; 79:1419–25.

- 29. Christodoulides N, Nikolidakis D, Chondros P, et al. Photodynamic therapy as an adjunct to nonsurgical periodontal treatment: a randomized, controlled clinical trial. J Periodontol 2008; 79:1638-44.
- 30. López NJ, Socransky SS, Da Silva I, et al. Effects of metronidazole plus amoxicillin as the only therapy on the microbiological and clinical parameters of untreated chronic periodontitis. J Clin Periodontol 2006; 33:648-60.
- 31. Bernimoulin JP. Recent concepts in plaque formation. J Clin Periodontol 2003; 30:7-9.