

Prevalence of Lower and Upper Canines Mal Occlusion among Dental Colleges Students

Emad Hammody Abdulla

Department of Dentistry, IBN Sina University of Medical and Pharmaceutical Sciences, Baghdad, Iraq

ABSTRACT

Background: Canine impaction is known to be one of the most common abnormalities in the teeth. The aim of this study was to evaluate the prevalence of mal occlusion canines and canine teeth, among Iraqi dental students in Baghdad.

Methods: This study included 260 students (161 (61.9%), female and 99 participants (38.07% male) between the ages of 18 and 24, faculty of dentistry at Ibn Sina University of Medical and Pharmaceutical Sciences, Iraq, Baghdad from February 2021 to June 2022. The tooth was considered affected when it was not aligned with the rest of the teeth in any of the tooth braces. Data on age, sex, the number, position, localization (right/left) of teeth, number of affected teeth, arch in question and tooth name were obtained the clinical of patients.

Results: The results of the cases showed a total (11) of impacted mandibular canine teeth were found, with prevalence (4.2%) of impaction. The results of the cases showed a total (6) of impacted maxillary canine teeth were found, with prevalence (2.3%) of impaction. The total of (17) cases malocclusion in the sample with prevalence (6.5%).

Conclusions: Early detection of impacted teeth is critical to successful treatment and thus demographic studies are important. Although larger samples are needed, this study provides a basis as to the frequency and type of canines affected in this particular group.

Key words: Impacted, Malocclusion, Upper canine, Lower canine

HOW TO CITE THIS ARTICLE: Emad Hammody Abdulla, Prevalence of Lower and Upper Canines Mal Occlusion among Dental Colleges Students, J Res Med Dent Sci, 2023, 11 (01): 000-000

Corresponding author: Dr. Emad Hammody Abdulla

E-mail: emad.g@ibnsina.edu.iq

Received: 12-Dec-2022, Manuscript No. JRMDS-23-81977;

Editor assigned: 15-Dec-2022, PreQC No. JRMDS-23-81977 (PQ);

Reviewed: 29-Dec-2022, QC No. JRMDS-23-81977;

Revised: 13-Jan-2023, Manuscript No. JRMDS-23-81977 (R);

Published: 20-Jan-2023

INTRODUCTION

Dental anomalies containing alterations in number, size and teeth structure often outline a major challenge for dental practitioners [1]. The affected tooth is defined as the tooth retained after the average time of its eruption [2]. It maintained that when tooth is unexploded after more than one year of natural age of eruption, is defined as "impacted" [3]. Tooth impaction is defined as "a condition in which a tooth is prevented from erupting via a physical barrier in a track of eruption" [4]. Next to third molar, upper jaw canine is the second most affected tooth in tooth arch [5,6]. Most affected teeth by impaction were reported as, third molars, maxillary canines, maxillary central incisors and premolars. These variable measures may have occurred using various age groups, sample sizes and patient variety [7-10]. The spread of affected teeth varies in proportion to the population, then be situated to range from (6.9%-76.6%) [11-15]. It's a recurring phenomenon has been extensively reported in the

literature [16-19]. Though, there are differences in spread of affected teeth in diverse parts of the world and their scattering in the upper in addition to lower jaws [20-22]. It was also informed that prevalence of affected teeth was more recurrent in females than in males. Furthermore, different factors attendant with dental curvature that includes different oldness groups, timing of the emergence of teeth, race or areas of study members and radial evaluation standards [23]. Impaction of maxillary canine can raise risk of root extrusion of neighboring lateral incisors, gum infections and cystic follicular lesions [24]. Canine impaction recognized to be one of the most common abnormalities in the teeth [25]. After the third molars were affected, permanent maxillary canines were furthestmost affected one [26]. Incidence of maxillary canine impaction was (20 times) higher than that of lower canine impaction [27]. The canines were impacted with the man to thousands years. A case reported in engraved skull old (2700-2724) BC, so it looks likely that an anomalous eruption of man Teeth are not caused by modified environments created by recent civilization [28]. Permanent canines the foundation and the column of the aesthetic and well-designed smile occlusion. Standing in corner of the teeth arch shaping the protrusion of canine for support from base of the alar and the upper lip. Functionally; It maintenances teeth that contribute to

DISCUSSION

In the study by Raja A Mustafa and Amal H Abuaffan, the proportion of impacted permanent canine was institute to be 2% [31]. Large figures of studies have shown the prevalence of permanently affected canine in various populations. Permanent canine among different populations, in upper jaw stated between them 0.8% and 5.4% [32-37]. While a smaller percentage has been reported in the lower jaw 0.3% and 0.44%. The results among Swedish and Sudanese population were similar (2.05%) [38]. Yet, it was less than the score between Saudi population (3.6%) and Turkish (3.73%) [39]. Prevalence of maxillary impaction between Sudanese was (1.6%) same (1.7%) in Swedish and Colombian children. A higher prevalence (3.29%) reported in Turkish patients and (5.4%) in Hungarian. Less one (0.8%) reported in Chinese patients [40]. The Sudanese mandibular canine impaction study (0.33%), a Swedish result, was relatively similar (0.35%). While he had a higher rate (0.44%) among Turkish patients. The difference among maxillary and mandibular impaction can be recognized to fact that the maxillary canines are the last teeth to grow and move a long path before the dental arch erupts, aggregate the potential for mechanical disorders leading to displacement besides impaction [41].

Higher prevalence of impaction can be accredited to the younger age collection of patients as well as to the ethnic differences between them. In the Sudanese study, most students (73.5%) had a solo effect, 24.5% had two canines and the lowest was (22.9%) informed in the Saudi population. On other hand, the Swedish noted only (8%) of the effects of bilateral canines. The male to female ratio in affected canines was 1:4 in the Sudanese study, in the Hungarian the ratio was 46:55, while in Turkish 1:1.51. Beck stated that the difference between the sexes credited to biological phenomena with genetic links, involving sex chromosomes. Abdulaziz Alamri, showed in his study that there were no significant variances in incidence of impacted teeth between male and female study members, which is in agreement with the results of other studies [42-48].

Exploration of radiographs revealed that prevalence of impacted teeth was (13.2%) in Saudi patients, close to other studies, for example 13.7% among the Greek population, 14.1% studied in the finland population and 16.8% in the northern part of India of the participants were diagnosed with impacted teeth, 44.1% in central part of Iran, 28.3% in China, In Hong Kong and Turkey, 3.58% of patients detected with canine impaction next 4,500 consecutive panoramic review.

Ahmed and Dhiaa the prevalence of misplaced canines is 11%, however, misplaced canines in the lower arch were higher (12%) than in the upper arch (10%) and the

distribution of misplaced canines was higher in males (11.3) compared to females (10.7) [49]. Iraqi studies show high prevalence in the lower arch compared to the upper arch for both sexes [50-53]. On the other hand Al-Huwaizi, et al. who found that impacted canines were more prevalent in the upper arch than in the lower arch [54]. Ahmed and Dhiaa, Aldabagh, et al. and Aldabagh found that canine malposition was higher in males than females [54,55].

In our study the prevalence of malposed canine is (6.5%) less than the results in previous similar Iraqi studies, but close to that in Saudi and Turkish one, the high prevalence of lower canine impaction (5%) than the upper one (1.5%) supported the Iraqi studies done in the past.

The prevalence in our study in female students is higher than male with no significant differences, This not agree with Iraqi studies but close to the ratio results in other countries.

There is need for large group students study includes many colleges of dentistry in many Universities, a survey like that carried in India and Arabia Saudi and other countries will be of a great help to fix the end view of the truth.

CONCLUSION

Early detection of impacted teeth is critical to successful treatment and thus demographic studies are important. Although larger samples are needed, this study provides a basis as to the frequency and type of canines affected in this particular group.

REFERENCES

1. Stecker SS, Beiraghi S, Hodges JS, et al. Prevalence of dental anomalies in a Southeast Asian population in the Minneapolis/saint Paul metropolitan area. *Northwest Dent* 2007; 86:25-28.
2. Pechalova PF, Pavlov NV, Konstantinova DA. Mandibular third molars in Southern Bulgaria: A clinical and radiological study of 1518 patients. *J Dent Probl Solut* 2017; 4:26-30.
3. Torres Lagares D, Flores Ruiz R, Infante Cossio P, et al. Transmigration of impacted lower canine. Case report and review of the literature. *Med Oral Patol Oral Cir Bucal* 2006; 11:171-174.
4. Rajendran R. *Shafer's textbook of oral pathology*. Elsevier India, Amsterdam, Netherlands, 2009.
5. Shah RM, Boyd MA, Vakil TF. Studies of permanent tooth anomalies in 7886 canadian individuals. *J Dent* 1978; 44:262-264.
6. Murray P, Brown NL. The conservative approach to managing unerupted lower premolars two case reports. *Int J Paediatr Dent* 2003; 13:198-203.

7. Lee PP. Impacted premolars. *Dent Update* 2005; 32:152-154.
8. Karapanou V. Endodontic management of an impacted premolar. *J Clin Pediatr Dent* 2005; 29:293-298.
9. McNamara C, McNamara TG. Mandibular premolar impaction: 2 case reports. *J Can Dent Assoc* 2005; 71:859-863.
10. Chu FCS, Li TKL, Lui VKB, et al. Prevalence of impacted teeth and associated pathologies a radiographic study of the Hong Kong Chinese population. *Hong Kong Med J* 2003; 9:158-163.
11. Hattab FN, Rawashdeh MA, Fahmy MS. Impaction status of third molars in Jordanian students. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1995; 79:24-29.
12. Peltola JS: A panorama tomographic study of the teeth and jaws of finnish university students. *Community Dent Oral Epidemiol* 1993; 21:36-39.
13. Yamaoka M, Furusawa K, Yamamoto M. Influence of adjacent teeth on impacted third molars in the upper and lower jaws. *Aust Dent J* 1995; 40:233-235.
14. Torres Lagares D, Flores Ruiz R, Infante Cossio P, et al. Transmigration of impacted lower canine. *Med Oral Patol Oral Cir Bucal* 2006; 11:171-174.
15. Shah RM, Boyd MA, Vakil TF. Studies of permanent tooth anomalies in 7886 canadian individuals. Impacted teeth. *Dent J* 1978; 44:262-264.
16. Aitasalo K, Lehtinen R, Oksala E. An orthopan tomography study of prevalence of impacted teeth. *Int J Oral Surg* 1972; 1:117-120.
17. Ahlqwist M, Grondahl HG. Prevalence of impacted teeth and associated pathology in middle aged and older Swedish women. *Community Dent Oral Epidemiol* 1991; 19:116-119.
18. Brown LH, Berkman S, Cohen D, et al. A radiological study of the frequency and distribution of impacted teeth. *J Dent Assoc S Afr* 1982; 37:627-630.
19. Peltola JS. A panorama tomographic study of the teeth and jaws of Finnish university students. *Community Dent Oral Epidemiol* 1993; 21:36-39.
20. Fardi A, Kondylidou Sidira A, Bachour Z, et al. Incidence of impacted and supernumerary teeth: A radiographic study in a North Greek population. *Medicina Oral Patologia Oral Y Cirugia Bucal* 2011; 16:e56- e61.
21. Chu FC, Li TK, Lui VK, et al. Prevalence of impacted teeth and associated pathologies: A radiographic study of the Hong Kong Chinese population. *Hong Kong Med J* 2003; 9:158-163.
22. Afify AR, Zawawi KH. The prevalence of dental anomalies in the Western region of Saudi Arabia. *ISRN Dentistry* 2012; 5.
23. Borges AH, Pedro FLM, Band´eca MC, et al. Prevalence of impacted teeth in a Brazilian subpopulation. *J Contemp Dent Pract* 2014; 15:209-213.
24. Richardson G, Russell KA. A review of impacted permanent maxillary cuspids diagnosis and prevention, *Journal (Canadian Dental Association)* 2000; 66:497-502.
25. Patil S, Maheshwari S. Prevalence of impacted and supernumerary teeth in the North Indian population. *J Clin Exp Dent* 2014; 6:e116-e120.
26. Becker A, Chaushu S. Etiology of maxillary canine impaction: A review. *American J Orthod Dentofacial Orthop* 2015; 148:557-567.
27. Kuftinec MM, Shapira Y. The impacted maxillary canine (II). orthodontic considerations and management. *Quintessence Int Dental Digest* 1984; 15:921-926.
28. Rajic S, Muretic Z, Percac S. Impacted canine in a prehistoric skull. *Angle Orthod* 1996; 66:477-480.
29. Mulick JF. Impacted Canines. *JC Orthod* 1997; 40:824-834.
30. Moyers RE. *Handbook of Orthodontics*. 4th edition, New York: Mosby, *Handbook of Orthodontics*. 1973; 55-58.
31. Mustafa RA, Abuaffan AH. Mustafa R, Abuaffan A. Prevalence of impacted canines among sudanese university students. *Braz Dent Sci* 2014; 17:27-33.
32. Dewel BF. The upper cuspid: Its development and impaction. *Angle Orthod* 1949; 19:79-90.
33. McSherry PF. The ectopic maxillary canine: A review. *Brit J orthod* 1998; 25:209-216.
34. Zahrani AA. Impacted cuspids in a Saudi population: Prevalence, etiology and complication. *Egypt Dent J* 1993; 39:367-374.
35. Rozsa N, Fabian G, Szadeczky B, et al. Prevalence of impacted permanent upper canine and its treatment in 11-18 year-old orthodontic patients. *Fogorv Sz* 2003; 96:65-69.
36. Peck S, Peck L, Kataja M. The palatally displaced canine as a dental anomaly of genetic origin. *Angle Orthod* 1994; 64:249-256.
37. Kokich VG, Mathews DP. Surgical and orthodontic management of impacted teeth. *Dent Clin North Am* 1993; 37:181-204.
38. Ericson S, Kurol J. Radiographic assessment of maxillary canine Eruption in children with clinical signs of eruption disturbances. *Eur J Orthod* 1986; 8:133-140.
39. Aydin U, Yilmaz HH, Yildirim D. Incidence of canine impaction and transmigration in a patient population. *Dentomaxillofac Radiol* 2004; 33:164-169.
40. Chu FC, Li TK, Lui VK, et al. Prevalence of impacted teeth and associated pathologies a radiographic study of Hong Kong Chinese population. *Hong Kong Med J* 2003; 9:158-163.
41. McSherry P, Richardson A. Ectopic eruption of the maxillary canine Quantified in three dimensions

- on cephalometric radiographs between The ages of 5 and 15 years. *Eur J Orthod* 1999; 21:41-48.
42. Alamri A, Alshahrani N, Abdullah Al-Madani, et al. Research article prevalence of impacted teeth in Saudi patients attending dental clinics in the eastern province of Saudi Arabia: A radiographic retrospective study. *Sci World J* 2020; 6.
 43. Fardi A, Kondylidou Sidira A, Bachour Z, et al. Incidence of impacted and supernumerary teeth: A radiographic study in a North Greek population. *Med Oral Patol Oral Cir Bucal* 2011; 16:e56- e61.
 44. Chu FC, Li TK, Lui VK, et al. Prevalence of impacted teeth and associated pathologies: A radiographic study of the Hong Kong Chinese population. *Hong Kong Med J* 2003; 9:158-163.
 45. Afify AR, Zawawi KH. The prevalence of dental anomalies in the Western region of Saudi Arabia. *ISRN Dent* 2012; 2012:5.
 46. Borges AH, Pedro FLM, Bandeca MC, et al. Prevalence of impacted teeth in a Brazilian subpopulation. *J Contemp Dent Pract* 2014; 15:209-213.
 47. Kramer RM, Williams AC. The incidence of impacted teeth: A survey at Harlem Hospital. *Oral Surg Oral Med Oral Pathol* 1970; 29:237-241.
 48. Arabion H, Gholami M, Dehghan H, et al. Prevalence of impacted teeth among young adults: A retrospective radiographic study. *J Dent Mater Tech* 2017; 6:131-137.
 49. Yaseen AA, Aldabagh DJ. Canine malposition in Iraqi dental students. *J Bagh College Dent* 2020; 32:16-18.
 50. Abdulla NM. Occlusal features and perception: A sample of 13-17 years old adolescent. Master Thesis, College of Dentistry, Baghdad University, Iraq, 1996.
 51. Batayina FAM. Occlusal features and perception of occlusion of Jordanian adolescents: A comparative study with an Iraqi sample. Master Thesis, College of Dentistry, University of Baghdad, Iraq, 1997.
 52. Al-Huwaizi AF, Al-Mulla AA, Al-Alousi WS. Dental crowding or spacing in 13 year olds. *J Coll Dent* 2002; 14:126-136.
 53. Aldabagh DJ, Ali FA, Al-Huwaizi AF. Buccally malposed canines in Yemen. *J Coll Dent* 2003; 15:18-21.
 54. Al-Huwaizi AF, Al-Mulla AA, Al-Alousi WS. Anomalies of tooth eruption and development in Iraqi 13 year olds a national survey. *J Coll Dent* 2002; 14:48-60.
 55. Aldabagh DJ. Buccally malposed mandibular canine in Iraqi Kurdish population. *Iraqi Orthod J* 2011; 7:6-9.