

A Retrospective Analysis of Prevalence of Anterior Open Bite in Patients Reporting to a Teaching Hospital in Chennai

Remmiya Mary Varghese*, Azima Hanin SM

Department of Orthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai

ABSTRACT

Introduction: Anterior Open Bite (AOB) is a debilitating malocclusion that requires thorough diagnosis to ensure long-term stability and proper treatment. The aim of this study was to determine the prevalence of AOB in patients visiting a university teaching hospital in Chennai, India in order to assess the need for orthodontic treatment and to determine its relation to other associated features.

Material and methods: This retrospective study was conducted among patients who were diagnosed with anterior open bite in a university teaching hospital in Chennai during the period of December 2020 to May 2021. The collected data was then subjected to statistical analysis using Statistical Package for Social Science (SPSS). Descriptive statistics and Chi square tests were used.

Results: Anterior open bite was frequently observed in males with 52.4% of the total population. Patients with anterior open bite had class I malocclusion prevalently with 85.1% of the total population. 89% of patients with anterior open bite had competent lips, followed by 6.8% of patients who had potentially competent lips. It was noticed that potentially competent lips were more frequent with females than in males.

Conclusion: AOB was more prevalent in males compared to females. Patients diagnosed with anterior open bite predominantly had Class I malocclusion and competent lips. Careful diagnosis and treatment plan for AOB should be done as any error in identifying the etiology may lead to a poor result. Due to the close connection between the prevalence of anterior open bite and dysfunctional problems, prevention modalities should be integrated into a national public health program. Awareness programs targeting parents should be implemented to reduce the risk of anterior open bite malocclusion.

Key words: Anterior open bite, Debilitating malocclusion, Diagnosis, Orthodontic treatment

HOW TO CITE THIS ARTICLE: Remmiya Mary Varghese, Azima Hanin SM, A Retrospective Analysis of Prevalence of Anterior Open Bite in Patients Reporting to a Teaching Hospital in Chennai, J Res Med Dent Sci, 2022, 10 (5): 068-076.

Corresponding author: Remmiya Mary Varghese

E-mail: remmiyav.sdc@saveetha.com

Received: 21-Feb-2022, Manuscript No. JRMDS-22-P-47963;

Editor assigned: 23-Feb-2022, Pre QC No. JRMDS-22-P-47963 (PQ);

Reviewed: 9-Mar-2022, QC No. JRMDS-22-P-47963;

Revised: 22-Apr-2022, Manuscript No. JRMDS-22-P-47963 (R);

Published: 2-May-2022

INTRODUCTION

Malocclusions may be considered a major public health problem due to the high rates of prevalence and treatment needs and the social impact these conditions cause [1]. Knowledge of etiology of malocclusion is extremely essential for the success of orthodontic treatment, since eliminating the cause is a prerequisite for correction of any malocclusion. In view of the increasing interest in early diagnosis and corresponding emphasis on preventive procedures, further information on etiological factors which cause malocclusion is needed [2].

The term open bite refers to no contact between anterior or posterior teeth. In the year 1842, Caravelli coined the

term "open bite" as a distinct classification of malocclusion which can be defined in different manners without any specificity. The term open bite is referred to as no opposing contact between anterior or posterior teeth. The complexity of open bite is attributed mainly to a combination of skeletal, dental and habitual factors. In recent literature, some authors have determined that open bite, or a tendency toward open bite, may occur due to smaller openbite than the normal. While one school of thought argues that open bite is characterized by end-on incisal relationships, the others state that there is no incisal contact present before diagnosing open bite [3-7]. Open bite was also defined by two other authors Sakuda and Subtelny [8]. As the open vertical dimension between the incisal edges of the maxillary and mandibular anterior teeth, although the deficiency in vertical dimension can occur between the anterior or the buccal segment.

Anterior open bite and posterior crossbite are one of the most prevalent debilitating malocclusions [9,10]. Anterior open bite develops at an early age, but might self-correct

spontaneously as the child grows. Malocclusion is the result of a combination of both etiological factored genetic and environmental [11]. Non-nutritive sucking habits are one of the main etiological factors correlated with anterior open bite. While literature states that anterior open bite is associated with non-nutritive sucking habits and other environmental factors with [12-14]. Few investigations have sought to establish associations between sociodemographic factors and malocclusion especially in children aged 5 years or less [9]. Moreover, most studies analyse predisposing factors separately, without considering the concurrent impact or assessing potential interactions. Clinical interest in the etiological factors and primitive diagnosis of malocclusion justifies epidemiological investigations focused on the main types of malocclusion found in preschool children.

Worldwide, the prevalence of anterior open bite in previous literature ranged between 1.6% and 47.1% [15-18]. Anterior open bite was diagnosed more frequently in children than in adults (1.5-24.5%) [17,19]. The treatment of open bite still remains a tough challenge to the clinician; careful diagnosis and timely intervention with proper treatment modalities and appliance selection will improve the treatment outcomes and long-term stability [20-22]. For complicated cases orthognathic surgery is the last and only resort [23]. However, the tendency toward relapse after conventional or surgical orthodontic treatment has been proved and stated in literature. Therefore, open bite is considered one of the most challenging dentofacial deformities to treat. There is a lack of literature about anterior open bite pertaining to the south Indian population. Our team has extensive knowledge and research experience that has translated into high quality publications [24-39]. The main aim of this study was to determine the prevalence of AOB in patients visiting a university teaching hospital in Chennai, India in order to assess the need for orthodontic treatment and to determine its relation to other associated features.

MATERIALS AND METHODS

Study setting: This university hospital-based retrospective study was carried out by reviewing the dental records of patients who underwent surgical extraction of impacted premolars who had visited a university teaching hospital in Chennai. Since this was a university hospital setting the large sample size and distribution of population contributed a major advantage for this study. Data collected was reliable and with evidence. The study was conducted after obtaining approval from the Institutional Ethical Review Board.

Sampling: Data was reviewed and collected from 86,000 patient records over a six months period from December 2020 to May 2021. Data of those patients who underwent extraction of impacted premolar was collected. From

2732 patients who were diagnosed with an open bite, patients in the age group of 10-60 years, and anterior open bite were included in the study while those with incomplete hospital records were excluded from the study. Cross verification was done using photographs and radiographs.

Data collection: The following patient data were recorded as follows: hospital record number, gender, age, radiographic/dental diagnosis. The Total population of patients who were diagnosed with anterior open bite was 1609. Data collected was then exported to Microsoft Excel 2010.

Data analytics: The acquired data was subjected to statistical analysis. Microsoft Excel 2010 data spreadsheet was used for tabulation of parameters and later exported to the Statistical Package for Social Science (SPSS version 20.0) for Windows. Descriptive statistics were applied to the data and chi-square tests were applied at a level of significance of 5% ($P < 0.05$).

RESULTS

Anterior open bite was most frequently seen in males with 52.4% of the total population. Patients with anterior open bite had class I malocclusion prevalently with 85.1% of the total population. 89% of patients with anterior open bite had competent lips, followed by 6.8% of patients who had potentially competent lips. It was noticed that potentially competent lips were more frequent with females than in males (Figures 1 and 2).




Figure 1: This bar graph represents the lip competency present in patients diagnosed with anterior open bite. X-axis represents lip competency and the y-axis represents the percentage of patients. 89% of patients with anterior open bite had competent lips, followed by 6.8% of patients who had potentially competent lips.




Figure 2: Bar graph depicting association between lip competency of patients with anterior open bite and gender of the patients. X-axis represents the gender with lip competency in the X cluster and Y-axis represents the number of patients who were diagnosed with anterior open bite. Blue colour denotes competent lip, green colour denotes incompetent lips and beige denotes potentially competent lips. *Chi-square test* was done and the association was found to be statistically not significant. Pearson *chi square* value: 6.344; df: 9; p value: 0.705 (<0.05), hence statistically not significant proving there was no significant association between lip competency in anterior open bite patients and their gender. However, it was noticed that potentially competent lips were more frequent with females than in males.

DISCUSSION

From this study, it was observed that anterior open bite was most frequently seen in males with 52.4% of the total population. This is consistent with the results reported by Hameedullah with a low male to female ratio 2:1 of anterior open bite [18]. Moreover, in another study 1585 Yemeni university students' males were recorded to have more open bite than females with a ratio of 4:1 [28]. However, in contrast to this study's findings, numerous previous studies found that anterior open bite was more common in females than males [10,13,15-18]. Variations in the prevalence of anterior open bite among genders can be partially attributed to the fact that Indian parents are more concerned about the appearance of boys than girls and try to prevent and treat all kinds of the abnormalities concerning the teeth and unpleasant appearance on the smile of their sons at an early age.

Patients with anterior open bite had class I malocclusion prevalently with 85.1% of the total population which is in correlation with the study by Rohit [40]. Which stated that complex open bite is frequently associated with class-I and class-II malocclusions and rarely with class III malocclusion. The reason for this result may be the fact that the most common aetiological factor of anterior

open bite is habitual; like thumb sucking, pacifier use, mouth breathing, lips and tongue habits. Class II, III malocclusions are found to be evident due to hereditary reasons, hence the disparity in the results.

Lip incompetence is a characteristic clinical feature in patients diagnosed with anterior open bite [23]. However, it was observed in this study that 89% of patients with anterior open bite had competent lips, followed by 6.8% of patients who had potentially competent lips and it was noticed that potentially competent lips were more frequent with females than in males.

CONCLUSION

Open bite is the most difficult malocclusion to treat in orthodontic practice. Treatment modalities include functional appliances in growing children and surgeries in adults. Relapse rates are highest in this type of malocclusion and hence the difficulty in treating these patients. Anterior open bite was more prevalent in males than females. Patients diagnosed with anterior open bite predominantly had class I malocclusion and competent lips. Careful diagnosis and treatment plan for anterior open bite should be done as any error in identifying the etiology may lead to a poor result. Due to the close connection between the prevalence of anterior open bite and dysfunctional problems, prevention modalities should be integrated into a national public health program. Awareness programs targeting parents should be implemented to reduce the risk of anterior open bite malocclusion.

ACKNOWLEDGEMENTS

This research was done under the supervision of the Department of Orthodontics, Saveetha dental College and Hospital. We sincerely show gratitude to the corresponding guide who provided insight and expertise that greatly assisted the research.

CONFLICT OF INTEREST

There was no potential conflict of interest.

SOURCE OF FUNDING

The present project is supported by

- Saveetha Institute of Medical and Technical Sciences
- Saveetha Dental College & Hospitals
- Saveetha University
- Al Hassain Trading, KSA

REFERENCES

1. Peres KG, Traebert ES de A, Marcenes W. Differences between normative criteria and self-perception in the assessment of malocclusion. *Rev Saude Publica* 2002; 36:230-236.
2. Macena MCB, Katz CRT, Rosenblatt A. Prevalence of a posterior cross bite and sucking habits in

- Brazilian children aged 18-59 months. *Eur J Orthod* 2009; 31:357-361.
3. Artese A, Drummond S, do Nascimento JM, et al. Criterios para o diagnóstico e tratamento estável da mordida aberta anterior. *Dental Press J Orthod* 2011; 16:136-161.
 4. Pithon MM. Angle Class I malocclusion with anterior open bite treated with extraction of permanent teeth. *Dental Press J Orthod* 2013; 18:133-140.
 5. Cabrera M de C, Cabrera CAG, de Freitas KMS, et al. Lateral open bite: treatment and stability. *Am J Orthod Dentofacial Orthop* 2010; 137:701-711.
 6. Shapiro PA. Stability of open bite treatment. *Am J Orthod Dentofacial Orthop* 2002; 121:566-568.
 7. Cozza P, Mucedero M, Baccetti T, et al. Early orthodontic treatment of skeletal open-bite malocclusion: a systematic review. *Angle Orthod* 2005; 75:707-713.
 8. Subtelny JD, Daniel Subtelny J, Sakuda M. Open-bite: Diagnosis and treatment. *Am J Orthod*. 1964; 50:337-358.
 9. Peres KG, Barros AJD, Peres MA, et al. Effects of breastfeeding and sucking habits on malocclusion in a birth cohort study. *Revista de Saúde Pública*. 2007; 41:343-350.
 10. Melink S, Vagner MV, Hocevar-Boltezar I, et al. Posterior crossbite in the deciduous dentition period, its relation with sucking habits, irregular orofacial functions, and otolaryngological findings. *Am J Orthod Dentofacial Orthop* 2010; 138:32-40.
 11. Berneburg M, Zeyher C, Merkle T, et al. Orthodontic findings in 4-to 6-year-old kindergarten children from southwest Germany. *J Orofac Orthop* 2010; 71:174-186.
 12. Heimer MV, Tornisiello Katz CR, Rosenblatt A. Non-nutritive sucking habits, dental malocclusions, and facial morphology in Brazilian children: a longitudinal study. *Eur J Orthod* 2008; 38:580-585.
 13. Katz CRT, Rosenblatt A, Gondim PPC. Nonnutritive sucking habits in Brazilian children: effects on deciduous dentition and relationship with facial morphology. *Am J Orthod Dentofacial Orthop* 2004; 126:53-57.
 14. Corrêa-Faria P, Ramos-Jorge ML, Martins-Júnior PA, et al. Malocclusion in preschool children: prevalence and determinant factors. *Eur Arch Paediatr Dent* 2014; 15:89-96.
 15. Ize-Iyamu IN, Isiekwe MC. Prevalence and factors associated with anterior open bite in 2 to 5 year old children in Benin City, Nigeria. *Afr Health Sci* 2012; 12:446-451.
 16. El-Mesbahy B, ElShiekh M, Hanafy R. Prevalence of Anterior Open Bite and Its Etiological Factors among a Group of Egyptian Children: A Cross Sectional Study. *Egypt Dent J* 2021; 67:1871-1878.
 17. Cozza P, Baccetti T, Franchi L, et al. Sucking habits and facial hyperdivergency as risk factors for anterior open bite in the mixed dentition. *Am J Orthod Dentofac Orthop* 2005; 128:517-519.
 18. Urzal V, Braga AC, Ferreira AP. The prevalence of anterior open bite in Portuguese children during deciduous and mixed dentition Correlations for a prevention strategy. *Int Orthod* 2013; 11:93-103.
 19. Ravassipour DB, Powell CM, Phillips CL, et al. Variation in dental and skeletal open bite malocclusion in humans with amelogenesis imperfecta. *Arch Oral Biol* 2005; 50:611-623.
 20. Peres KG, De Oliveira Latorre MDRD, SHEIHAM A, et al. Social and biological early life influences on the prevalence of open bite in Brazilian 6-year-olds. *Int J Paediatr Dent* 2007; 17:41-49.
 21. Lopez-Perez R, Borges-Yanez SA, Lopez-Morales P. Anterior open bite and speech disorders in children with Down syndrome. *Angle Orthod* 2008; 78:221-227.
 22. Harila V, Heikkinen T, Gron M, et al. Open bite in prematurely born children. *J Dent Child* 2007; 74:165-170.
 23. Ngan P, Fields HW. Open bite: a review of etiology and management. *Pediatr Dent* 1997; 19:91-98.
 24. Felicita AS. Orthodontic extrusion of Ellis Class VIII fracture of maxillary lateral incisor. The sling shot method. *Saudi Dent J*. 2018;30:265-269.
 25. Arvind P TR, Jain RK. Skeletally anchored forsus fatigue resistant device for correction of Class II malocclusions-A systematic review and meta-analysis. *Orthod Craniofac Res* 2021; 24:52-61.
 26. Chandrasekar R, Chandrasekhar S, Sundari KKS, et al. Development and validation of a formula for objective assessment of cervical vertebral bone age. *Prog Orthod* 2020; 21:38.
 27. Alam MK, Alfawzan AA, Haque S, et al. Sagittal Jaw Relationship of Different Types of Cleft and Non-cleft Individuals. *Front Pediatr* 2021; 9:651951.
 28. Marya A, Venugopal A. The Use of Technology in the Management of Orthodontic Treatment-Related Pain. *Pain Res Manag* 2021; 55:12031.
 29. Adel S, Zaher A, El Harouni N, et al. Robotic Applications in Orthodontics: Changing the Face of Contemporary Clinical Care. *Biomed Res Int* 2021; 9954615.
 30. Venugopal A, Vaid N, Bowman SJ. Outstanding yet redundant. After all, you may be another Choluteca Bridge. *Semin Orthod* 2021; 27:53-56.
 31. Gopalakrishnan U, Felicita AS, Mahendra L, et al. Assessing the Potential Association Between Microbes and Corrosion of Intra-Oral Metallic Alloy-Based Dental Appliances Through a Systematic Review of the Literature. *Front Bioeng Biotechnol* 2021; 9:154.
 32. Marya A, Karobari MI, Selvaraj S, et al. Risk Perception of SARS-CoV-2 Infection and Implementation of Various Protective Measures by Dentists Across Various Countries. *Int J Environ Res Public Health* 2021; 18.

33. Ramesh A, Varghese S, Jayakumar ND, et al. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients- A case control study. *J Periodontol* 2018; 89:1241-1248.
34. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol* 2021; 122:105030.
35. Joseph B, Prasanth CS. Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2021; 132:118-119.
36. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med* 2019; 48:115-121.
37. Duraisamy R, Krishnan CS, Ramasubramanian H, et al. A Compatibility of Non original Abutments with Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments. *Implant Dent* 2019; 28:289-295.
38. Gothandam K, Ganesan VS, Ayyasamy T, et al. Antioxidant potential of theaflavin ameliorates the activities of key enzymes of glucose metabolism in high fat diet and streptozotocin-induced diabetic rats. *Redox Rep* 2019; 24:41-50.
39. Varghese RM, Subramanian AK, Sreenivasagan S, et al. Comparison of dentoskeletal changes in skeletal class II cases using two different fixed functional appliances: Forsus fatigue resistant device and powerscope class II corrector—A clinical study. *J In Oral Health* 2021; 13:234.
40. Rohit K. Open bites malocclusion: An overview. *J oral health craniofac sci* 2018; 11-20.