

showed slightly alkaline pH as compared to normal individuals. The future study may be conducted with the broader representation of sample to verify the existing study finding.

REFERENCES

1. Maheswari TU. Assessment of salivary pH In patients with oral submucous fibrosis. Intern J Pharmaceu Res 2020; 12.
2. Panchbhai A. Effect of oral submucous fibrosis on jaw dimensions. Turkish J Orthodon 2019; 32:105.
3. Lalfamkima F, Bommasi S, Reddy K, et al. Comparative evaluation of alteration in salivary flow rate between betal nut/gutkha chewers with and without OSMF and healthy subjects: A prospective case-control study. Oncol J India 2021; 5:1.
4. Hande AH, Chaudhary MS, Gadmail AR, et al. Role of hypoxia in malignant transformation of oral submucous fibrosis. J Datta Meghe Instit Med Sci Univ 2018; 13:38-43.
5. Donoghue M, Basandi PS, Adarsh H, et al. Habit-associated salivary pH changes in oral submucous fibrosis-A controlled cross-sectional study. J Oral Maxillofacial Pathol 2015; 19:175.
6. Shih YH, Wang TH, Shieh TM, et al. Oral submucous fibrosis: A review on etiopathogenesis, diagnosis and therapy. Intern J Molec Sci 2019; 20:2940.
7. More CB, Rao NR, Hegde R, et al. Oral submucous fibrosis in children and adolescents: Analysis of 36 cases. J Indian Soc Pedodon Prev Dent 2020; 38:190-199.
8. Shih YH, Wang TH, Shieh TM, et al. Oral submucous fibrosis: A review on etiopathogenesis, diagnosis and therapy. Intern J Molec Sci 2019; 20:2940.
9. Sarode SC, Chaudhary M, Gadmail A, et al. Dysplastic features relevant to malignant transformation in atrophic epithelium of oral submucous fibrosis: A preliminary study. J Oral Pathol Med 2018; 47:410-416.
10. Shende V, Biviji AT, Akarte N. Estimation and correlative study of salivary nitrate and nitrite in tobacco related oral squamous carcinoma and submucous fibrosis. J Oral Maxillofacial Pathol 2013; 17:381.
11. Lee JM, Garon E, Wong DT. Salivary diagnostics. Orthodon Craniofacial Res 2009; 12:206-211.
12. Humphrey SP, Williamson RT. A review of saliva: Normal composition, flow and function. J Prosthe Dent 2001; 85:162-169.
13. Tamgadge P, Wasekar R, Kulkarni S, et al. Comparative evaluation of alteration in salivary pH among gutkha chewers with and without oral submucous fibrosis and healthy subjects: A prospective case-control study. J Curr Oncol 2020; 3:8.