

Suicidal Attempt in a Schizophrenic Patient with a High Level of Expressed Emotion in The Patient's Family

Rajab Bresaly^{1*}, Ammar Albokhari², Rotan Mal³, Abdulrahman Tashkandi³, Abdulrhman Alsawas³, Mohammed Abualshamat³, Ghassan Al Harazi³, Saeed Alamri⁸, Alaa Shabekni³, Ahmed Abdullah Mohsen Alharthi², Mari Osman Mari Alwadai², Maha Alluqmani³

¹Department of Psychiatry, Hera General Hospital, Makkah, Saudi Arabia

²Department of Psychiatry, Eradah Complex and Mental Health, Najran, Saudi Arabia

³Medical Resident, Hera General Hospital, Makkah, Saudi Arabia

ABSTRACT

Schizophrenia is a chronic mental disorder known by distortions in perception and thinking. Expressed emotion (EE) is an attitude towards a patient with a mental disorder from the individuals surrounding him. EE was predicted as a key-element in relapsing of schizophrenia through criticism, emotional over-involvement, and/or intrusiveness. EE has always been a known risk factor for relapsing in schizophrenia but in our case, that risk has increased the level of suicidology as few articles have linked EE as a risk factor to increase suicidality in other mental disorders. We present a case of young male from Saudi Arabia with a high level of expressed emotion from the family in the form of emotional over-involvement in better highlighting the importance of family therapy through psycho-educational interventions in managing of a schizophrenic patient.

Key words: Schizophrenia, Expressed Emotion, Suicide

HOW TO CITE THIS ARTICLE: Rajab Bresaly, Ammar Albokhari, Rotan Mal, Abdulrahman Tashkandi, Abdulrhman Alsawas, Mohammed Abualshamat, Ghassan Al Harazi, Saeed Alamri, Alaa Shabekni, Ahmed Abdullah Mohsen Alharthi, Mari Osman Mari Alwadai, Maha Alluqmani, Suicidal Attempt in a Schizophrenic Patient with a High Level of Expressed Emotion in The Patient's Family, J Res Med Dent Sci, 2022, 10 (11): 130-132.

Corresponding author: Rajab Bresaly

e-mail ✉: rbresaly@moh.gov.sa

Received: 24-Oct-2022, Manuscript No. JRMDs-22-78099;

Editor assigned: 26-Oct-2022, PreQC No. JRMDs-22-78099(PQ);

Reviewed: 09-Nov-2022, QC No. JRMDs-22-78099(Q);

Revised: 15-Nov-2022, Manuscript No. JRMDs-22-78099(R);

Published: 22-Nov-2022

CASE REPORT

An 18-year-old male patient arrived at the emergency department in a general hospital in Saudi Arabia with a deep cut wound of approximately 15 cm in length and multiple superficial cuts in the anterior aspect of his neck with heavy bleeding. His vital signs were as follows: temperature of 37°C, heart rate of 78 beats/min, blood pressure of 108/65 mmHg, respiratory rate of 20 breaths/min, and oxygen saturation of 99%. Two units of packed red blood cells were transfused, and he was rushed to the operating room for airway securing and the neck vessels repair with the performance of tracheostomy procedure. Toxicology results were unremarkable for any illicit substances except for the presence of benzodiazepine, which was one of his medications during his current intensive care unit admission.

The psychiatry team was consulted; upon the interview, the patient was exhibiting cognitive distortions like memory impairment and inattentive to communication. However, physical examination revealed no signs of scars, cutting wounds, needle bricks, and skin mutilation in both upper limbs. Collateral history was provided by his parents who reported that at the age of 17, their

INTRODUCTION

Schizophrenia is a chronic mental disorder characterized by distortions of perception and thinking [1]. Suicide is intentional attempt to end one's own life [2]. Since 1990, suicide has been one of the top 20 causes of death and it was ranked 15th in most recent data [3], accounting for 1.5% of mortality worldwide [4]. In 2019, Saudi Arabia had an estimated suicide rate of 5.4 per 100,000 populations according to the World Health Organization, and it was higher in males [5]. Expressed emotion (EE) is an attitude towards a patient with a mental disorder from the individuals surrounding him [6]. High EE was predicted as a key element in the relapse of schizophrenia through criticism, emotional over-involvement, and/or intrusiveness [7,8].

son was already hearing voices and had aggressive behavior toward the whole family to the point that his siblings will lock themselves during aggressive episodes. As the family strongly believed that it was a demonic possession or a consequence of witchcraft, they initially asked for the help of the local sheik. After weeks of trials with religious methods through multiple sheiks, they finally brought their son to the psychiatric emergency department for consult, which resulted in prescription medications without admission. The parents did not fully comprehend the severity of the mental disorder and the significance of medication adherence to a patient with psychosis, as they continued with the religious methods of treating their son. The psychiatry team recommended starting quetiapine 100 mg orally twice a day; alprazolam 0.5 mg orally once at bedtime; and escitalopram 20 mg orally once at bedtime, with urgent referral to the mental health hospital for admission after being discharged from his surgical condition. After two weeks of hospitalization with continued psychotropic medications, the patient was interviewed by the psychiatrist and exhibited normal behavioral and thought processing with negative psychotic symptoms and guilt feeling regarding the suicidal attempt. After his mother refused to admit him into the mental health hospital, the patient was discharged and allowed to continue his antipsychotic and Serotonin Selective Reuptake Inhibitor (SSRI) medications at home, with a scheduled one-week return visit to the psychiatry clinic; however, the patient did not show up.

Second admission

One month after his first admission, the patient's relatives brought him to the emergency department due to aggressive behavior two days before his presentation. Prior to his admission, his family needed a second opinion regarding his psychiatric condition; therefore, they sought consult with another psychiatrist, causing a complete change in his psychotropic medications from two to four medications, which further resulted from the initial decreased medication compliance to completely withholding treatment. This time, the family agreed for admission into the mental health hospital.

DISCUSSION

A study conducted in the Eastern region of Saudi Arabia reported that suicide by hanging was the most common method, accounting for approximately 90% of 126 cases, with one case of suicide by throat cutting (0.79%) [1]. A patient with schizophrenia has a 40.8% risk of suicidal attempt at least once in his lifetime, with that risk decreasing to a slightly lower percentage of 39.6% on recurrence [9]. One of the known risk factors for the relapse of patients with schizophrenia is the high level of EE in the patient's family [7,10]. In this study, the emotional over-involvement of the family by denying the presence of a mental disorder and holding into a superstitious belief of demonic possession or witchcraft caused a major delay in patient treatment.

Moreover, even after seeking consult from the psychiatry emergency department, the family remained in denial to start the psychotropic medications, resulting in the patient's aggressive form of suicide attempt. The high level of EE has a likelihood of increasing the probability of suicide in a patient with a mental disorder compared with a low level of EE [11,12]. Psych education, emotional processing, stress reduction, structured problem solving, and cognitive reappraisal are methods of family therapies that need to be delivered by a mental health professional [7]. Psych educational interventions must be used in family members and caregivers as early as possible in the first episode of psychosis [13,14].

CONCLUSION

The high level of EE in a patient's family can result in a devastating effect on the patient's well-being, as in this case. Therefore, high EE has a possible risk of increasing the likelihood of suicide; therefore, we urge our fellow psychiatrists to be keen in looking for high EE in their interviews and must use psycho educational interventions as early as possible and shift to more complex interventions thereafter.

ETHICS STATEMENT AND CONFLICT OF INTEREST DISCLOSURES

Human subjects: Consent was obtained. Conflicts of interest: In compliance with the ICMJE uniform disclosure form, all authors declare the following: Payment/services info: All authors have declared that no financial support was received from any organization for the submitted work. Financial relationships: All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. Other relationships: All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

REFERENCES

1. <https://www.nimh.nih.gov/health/topics/schizophrenia>
2. Madadin M, Mahmoud A, Alsowayigh K, et al. Suicide deaths in Dammam, Kingdom of Saudi Arabia: Retrospective study. *Egyptian J Forensic Sci* 2013; 3:39-43.
3. <https://ourworldindata.org/causes-of-death>.
4. Martínez-Alés G, Jiang T, Keyes KM, et al. The recent rise of suicide mortality in the United States. *Annu Rev Public Health* 2022; 43:99-116.
5. <https://apps.who.int/gho/data/node.main.MHSUICIDEASDR?lang=en>.
6. Amaresha AC, Venkatasubramanian G. Expressed emotion in schizophrenia: An overview. *Indian J Psychol Med* 2012; 34:12-20.
7. Caqueo-Úrizar A, Rus-Calafell M, Urzúa A, et al. The role

-
- of family therapy in the management of schizophrenia: challenges and solutions. *Neuropsychiatr Dis Treat* 2015; 11:145.
8. Butzlaff RL, Hooley JM. Expressed emotion and psychiatric relapse: A meta-analysis. *Arch Gen Psychiatry* 1998; 55:547-552.
 9. Aydın M, İlhan BÇ, Tekdemir R, et al. Suicide attempts and related factors in schizophrenia patients. *Saudi Med J* 2019; 40:475.
 10. Ng SM, Fung MH, Gao S. High level of expressed emotions in the family of people with schizophrenia: has a covert abusive behaviours component been overlooked?. *Heliyon* 2020; 6:e05441.
 11. Ellis AJ, Portnoff LC, Axelson DA, et al. Parental expressed emotion and suicidal ideation in adolescents with bipolar disorder. *Psychiatry Res* 2014; 216:213-216.
 12. Demir S. The relationship between expressed emotion and the probability of suicide among Turkish psychiatric outpatients: A descriptive cross-sectional survey. *Fam Community Health* 2018; 41:111-116.
 13. Fabri Cabral RR, Chaves AC. Multi-family group intervention in a programme for patients with first-episode psychosis: A Brazilian experience. *Int J Soc Psychiatry* 2010; 56:527-532.
 14. Barrio C, Yamada AM. Culturally based intervention development: The case of Latino families dealing with schizophrenia. *Res Soc Work Pract* 2010; 20:483-492.